

Modelling short-term energetic costs of sonar disturbance to cetaceans using high-resolution foraging data

Max F. Czapanskiy¹  | Matthew S. Savoca¹  | William T. Gough¹  | Paolo S. Segre¹  |
Danuta M. Wisniewska^{1,2}  | David E. Cade^{1,3}  | Jeremy A. Goldbogen¹ 

¹Hopkins Marine Station, Department of Biology, Stanford University, Pacific Grove, CA, USA

²Centre d'Etudes Biologiques de Chizé, CNRS-Université de La Rochelle, Villiers-en-Bois, France

³Institute of Marine Sciences, University of California, Santa Cruz, CA, USA

Correspondence

Max F. Czapanskiy
Email: maxczap@stanford.edu

Funding information

U.S. Office of Naval Research, Grant/Award Number: N00014-16-1-2477; Stanford University's William R. and Sara Hart Kimball Fellowship; Data Science Scholar Fellowship

Handling Editor: Verena Trenkel

Abstract

1. Anthropogenic noise is a pervasive and increasing source of disturbance to wildlife. Marine mammals exhibit behavioural and physiological responses to naval sonar and other sound sources. The lost foraging opportunities and elevated locomotor effort associated with sonar disturbance likely carry energetic costs, which may lead to population-level consequences.
2. We modelled the energetic costs associated with behavioural responses using (a) empirical datasets of cetacean feeding rates and prey characteristics and (b) allometry of swimming performance and metabolic rates.
3. We applied our model to compare the short-term (i.e. the scale of the disturbance response; hours to days) energetic costs of a variety of observed behavioural responses. Efficient foragers (e.g. baleen whales) incur a greater relative energetic cost for mild behavioural responses as compared to the most extreme observed response for larger odontocetes (e.g. beaked whales). Energetic costs are more sensitive to lost feeding opportunities than increased energy expenditure from elevated locomotor effort.
4. To scale up from short-term costs to long-term effects (months to years), future research should address individuals' capacity to compensate for energetic losses as well as energetic thresholds for demographic rates (survival, fecundity). We discuss how relative energetic costs correlate with species' pace of life and the implications for conservation planning.
5. *Synthesis and applications.* Current approaches towards understanding the Population Consequences of Disturbance (PCoD) often must rely on expert opinion due to data deficiency. Our model provides an empirical method for linking behaviour to energetics, which is critical for managers to make informed decisions on actions that may affect marine mammal species. Furthermore, our model is applicable to other forms of disturbance, such as vessel traffic or seismic exploration, and our scaling approach enables risk projections for understudied species.

Abbreviations: A, Stroke amplitude (m); C_L , Mass-specific locomotor cost of a stroke ($\text{kJ stroke}^{-1} \text{kg}^{-1}$); E, Energy cost of disturbance (kJ); E^* , Relative energetic cost of disturbance (unitless); E_p , Energy acquired per prey capture event (kJ/event); f, Stroke frequency (Hz); Δf , Change in stroke frequency (Hz); P_a , Rate of energy acquisition (kJ/hr); ΔP_e , Change in rate of energy expenditure (kJ/hr); PCoD, Population Consequences of Disturbance; r_p , Feeding rate (event/hr); St, Strouhal number (unitless); t_d , Time displaced from feeding (hr); t_r , Time spent in flight response (hr); U_p , Swim speed during flight response (m/s).

KEYWORDS

anthropogenic noise, cetaceans, disturbance, energetics, foraging behaviour, predator-prey, sonar, sublethal effects

1 | INTRODUCTION

Human activity is profoundly altering terrestrial and marine soundscapes, with impacts expected to increase in the future (Barber et al., 2010; Frisk, 2012; Kaplan & Solomon, 2016; McDonald et al., 2006). Like other forms of disturbance, anthropogenic noise can induce behavioural and physiological responses with population-level consequences (Frid & Dill, 2002; Shannon et al., 2016; Sutherland, 2007). Effects can be acute, including mortality (Simpson et al., 2016), or chronic, such as reduced reproductive success (Injaian et al., 2018; Kleist et al., 2018). Sound moves more efficiently in water than in air, so the impacts of increasing anthropogenic noise are particularly cause for concern in marine ecosystems (Duarte et al., 2021; Hildebrand, 2009; Popper & Hawkins, 2012).

Sound is a critical aspect of cetacean life history, especially for toothed whales (Odontoceti). Odontocetes (porpoises, dolphins, beaked and sperm whales), use echolocation to locate and capture prey in light-limited environments, from turbid rivers to ocean depths up to 3 km (Lindberg & Pyenson, 2007; Shearer et al., 2019). Both toothed and baleen whales (Mysticeti) use sound in communication and social behaviour (Jensen et al., 2012; Sørensen et al., 2018; Stafford et al., 1998; Tyack, 1986). Critically, in certain states and contexts, cetaceans may respond to anthropogenic sound

sources, sometimes with fatal results. Severe responses to military mid-frequency active sonar have been directly linked to mortality in odontocetes (Cox et al., 2006; D'Amico et al., 2009; Jepson et al., 2003; Parsons, 2017). Sonar-associated mass stranding events in the 1990s and 2000s led to a series of behavioural response studies (Table 1), showing that cetaceans exposed to sonar cease foraging, swim faster and flee the sound source (Harris et al., 2017; Southall et al., 2016). The biological significance of these sublethal responses remains largely unknown.

Behavioural responses, such as foraging cessation and elevated locomotion, carry energetic costs that may negatively impact individuals and populations. However, behavioural changes alone do not necessarily lead to population declines (Gill et al., 2001; Griffin et al., 2007). The Population Consequences of Disturbance (PCoD) conceptual framework addresses this issue using transfer functions to link changes in individuals' behaviour and physiology to population dynamics (Harwood et al., 2016; Pirodda et al., 2018). However, data are lacking for many of these transfer functions, necessitating expert solicitation for model parameterization (King et al., 2016). These 'interim' models have provided valuable insight by demonstrating, for example, how life-history stage (Farmer et al., 2018; Villegas-Amtmann et al., 2015) and resource availability (Hin et al., 2019; New et al., 2013) can mitigate or exacerbate the effects

TABLE 1 Controlled exposure experiments quantifying behavioural responses of cetaceans to low- and mid-frequency active sonar. Mysticetes in bold. N is the number of individuals exposed to sonar

Species	N	Region	Example feeding cessation	Example flight response	Reference
<i>Berardius bairdii</i>	1	E Pacific	50 min	30 min, 3 m/s	Stimpert et al. (2014)
<i>Globicephala melas</i>	6	Arctic			Antunes et al. (2014)
<i>Hyperoodon ampullatus</i>	1	Arctic	>8 hr		Miller et al. (2015)
<i>Hyperoodon ampullatus</i>	1	Arctic	>6 hr	10 min, 4 m/s	Sivle et al. (2015)
<i>Hyperoodon ampullatus</i>	3	Arctic	>8 hr	12 min, 3 m/s	Wensveen et al. (2019)
<i>Mesoplodon densirostris</i>	1	W Atlantic			Tyack et al. (2011)
<i>Orcinus orca</i>	8	Arctic			Miller et al. (2014)
<i>Ziphius cavirostris</i>	2	E Pacific	6.6 hr	1.6 hr, 2.6 m/s	DeRuiter et al. (2013)
<i>Balaenoptera acutorostrata</i>	1	Arctic			Sivle et al. (2015)
<i>Balaenoptera acutorostrata</i>	2	E Pacific, Arctic	70 min		Kvadsheim et al. (2017)
<i>Balaenoptera musculus</i>	17	E Pacific	62 min	5 min, 3 m/s	Goldbogen et al. (2013)
<i>Balaenoptera musculus</i>	9	E Pacific			Friedlaender et al. (2016)
<i>Balaenoptera musculus</i>	42	E Pacific	>1 hr	5 min, 3 m/s	Southall et al. (2019)
<i>Megaptera novaeangliae</i>	11	Arctic			Sivle et al. (2015)
<i>Megaptera novaeangliae</i>	13	Arctic			Sivle et al. (2016)

of disturbance, but they also highlight the need for empirically quantifying undisturbed behaviour, physiology and demography to establish a baseline.

Quantifying the foraging and movement energetics of wild cetaceans presents numerous challenges. Direct measurements of energy expenditure are limited to small- to medium-sized toothed whale species using respirometry or doubly labelled water techniques (Fahlman et al., 2016; Rojano-Doñate et al., 2018; Williams et al., 1993). Baleen whale metabolic rates have never been measured directly but have been inferred from models parameterized primarily with breathing rate data (Blix & Folkow, 1995; Sumich, 1983). For behavioural response studies, the theoretical relationship between exercise kinematics, such as fluke stroking rates, and swimming speed across body sizes (grounded in biomechanical constraints) provides a first approximation of increases in metabolism due to changes in locomotion (Dial et al., 2008; Fahlman et al., 2016). Stroking rates recorded for captive animals show a consistent relationship with speed and size (Rohr & Fish, 2004), which is supported by studies of wild animals (Gough et al., 2019; Sato et al., 2007; Watanabe et al., 2011, 2015).

Empirical estimates of energy intake are possible for a wider range of species, though still present logistical challenges. The biomass and energy obtained per feeding event for rorqual whales (the 'pleated' baleen whales such as blues, *Balaenoptera musculus*, and humpbacks, *Megaptera novaeangliae*) have been measured by combining active acoustics with animal-borne sensors and morphology of the engulfment apparatus (Goldbogen et al., 2019). Active acoustic techniques, also called prey mapping, use the backscatter of high-frequency sound from prey to measure the composition, extent and density of schools of crustaceans, squid and fish (Benoit-Bird & Lawson, 2016). Rorqual lunge feeding behaviour is characterized by distinct kinematic signatures that are readily captured by high-resolution animal-borne accelerometer tags. The whale accelerates as it approaches a prey school, engulfs a large quantity of prey-laden water and rapidly decelerates while filtering the engulfed water mass through baleen plates (Cade et al., 2016; Goldbogen et al., 2017). In contrast, toothed whales use echolocation to forage, and sound-recording tags can detect the acoustic signatures (i.e. high-rate click trains, or buzzes, and prey echoes) associated with prey capture (Johnson et al., 2004; Miller et al., 2004; Wisniewska et al., 2016). Odontocete diet composition and prey size have been estimated from non-digestible parts in stomach samples, such as squid beaks and fish otoliths (Clarke, 1996), and with active acoustic techniques (Wisniewska et al., 2016).

Here, we use empirical estimates of cetacean energy intake to predict the energetic costs of sonar disturbance and address the following questions. First, are energetic costs more sensitive to lost feeding opportunities (reduced acquisition) or elevated locomotor effort (increased expenditure)? We hypothesize that lost feeding opportunities are more important because of the low cost of cetacean locomotion (Williams, 1999) and high foraging efficiency conferred by adaptations such as echolocation and bulk filter feeding (Goldbogen et al., 2017; van der Hoop et al., 2019; Watwood

et al., 2006). Second, which species face the greatest immediate energetic costs relative to body size? The biological consequences of energy costs depend on energetic requirements; a 1,000 kJ cost will have greater consequences for a small porpoise than a large baleen whale. Mass-specific energy costs are inappropriate for comparison because energetic requirements (basal or field metabolic rate) are not isometrically proportional to body size (Kleiber, 1975; Nagy, 2005). Therefore, we define the relative energetic cost of a behavioural response as the ratio of its absolute energetic cost to the allometrically predicted daily basal metabolic rate. Although basal metabolic rates do not account for important energetic demands, such as lactation, they are correlated with many phylogenetic, physiological and ecological parameters and, as such, are a useful proxy for energy requirements (White & Seymour, 2004). This ratio facilitates interspecific comparisons by accounting for differential energy budgets across large body size ranges, but it is not appropriate for intraspecific comparisons where energy budget variability is influenced by other factors (such as life-history stage) more than body size. As feeding efficiency increases with body size among mysticetes, but decreases with body size among odontocetes (Goldbogen et al., 2019), we hypothesize that relative energetic costs are greatest for very large and small species compared to intermediate sizes. This scaling approach facilitates assessments for understudied species and is broadly applicable to other forms of disturbance, such as vessel traffic and seismic exploration.

2 | MATERIALS AND METHODS

We modelled the energetic cost of sonar exposure as the sum of energetic costs due to lost feeding opportunities and elevated locomotion during the flight response (Equation 1) for 11 species, selected on the basis of available feeding rate and prey energy content data (Table 2). See Figure 1 for a flowchart showing how the model combines species-specific energetics and behavioural scenarios to predict absolute and relative energetic costs.

$$E = P_a t_d + \Delta P_e (U_f) t_f, \quad (1)$$

where:

E is the energetic cost of sonar exposure in kJ.

P_a is the undisturbed rate of energy acquisition in kJ/hr.

t_d is the time displaced from feeding in hours.

ΔP_e is the increase in rate of energy expenditure when fleeing in kJ/hr (a function of swim speed).

U_f is the swim speed during the flight response in m/s.

t_f is the duration of the flight response in hours.

Together, t_d , U_f and t_f parameterize the behavioural response. Because the behavioural response can be highly variable, depending on the animal's behavioural state and the amplitude, frequency, and distance of the sound source, we leave these parameters to model users' discretion to allow greater flexibility (Ellison et al., 2012; Friedlaender et al., 2016). Energy acquisition and expenditure rates

TABLE 2 Cetacean species used in developing the model, covering three orders of magnitude in body size. Length and mass taken from (Goldbogen et al., 2019). Only *P. macrocephalus* exhibits strong sexual dimorphism; size here is for females. Mysticetes (all rorquals) in bold

Species	Length (m)	Mass (kg)	Family
Harbour porpoise (<i>Phocoena phocoena</i> , <i>Pp</i>)	1.2	31	Phocoenidae
Risso's dolphin (<i>Grampus griseus</i> , <i>Gg</i>)	3	350	Delphinidae
Blainville's beaked whale (<i>Mesoplodon densirostris</i> , <i>Md</i>)	4.1	860	Ziphiidae
Short-finned pilot whale (<i>Globicephala macrorhynchus</i> , <i>Gma</i>)	4.3	980	Delphinidae
Long-finned pilot whale (<i>Globicephala melas</i> , <i>Gme</i>)	5	1,200	Delphinidae
Cuvier's beaked whale (<i>Ziphius cavirostris</i> , <i>Zc</i>)	6.6	2,900	Ziphiidae
Minke whale (<i>Balaenoptera bonaerensis</i>, <i>Bb</i>)	7.8	6,700	Balaenopteridae
Sperm whale (<i>Physeter macrocephalus</i> , <i>Pm</i>)	11	15,000	Physeteridae
Humpback whale (<i>Megaptera novaeangliae</i>, <i>Mn</i>)	14	36,000	Balaenopteridae
Fin whale (<i>Balaenoptera physalus</i>, <i>Bp</i>)	20.2	53,000	Balaenopteridae
Blue whale (<i>Balaenoptera musculus</i>, <i>Bm</i>)	25.2	93,000	Balaenopteridae

are species-specific parameters based on empirical data and scaling relationships.

2.1 | Lost energy acquisition

The undisturbed rate of energy acquisition was modelled as the product of feeding rate and energy acquired per feeding event.

$$P_a = r_f E_p, \quad (2)$$

where:

r_f is the feeding rate in prey capture events/hr.

E_p is the energy acquired per prey capture event in kJ.

Feeding rates were estimated using kinematic and acoustic data from animal-borne bio-logging devices (Goldbogen et al., 2019). We excluded deployments involving controlled exposure experiments to get a best estimate of undisturbed feeding rates. Harbour porpoises




FIGURE 1 Flowchart explaining model components. Each species' energy intake (blue) and expenditure (red) parameters (first column) were estimated from empirical evidence or scaling relationships. The behavioural scenario (second column) quantifies the magnitude of the response to sonar. The energetic cost of the response (third column) is a function of the species-specific energetics mediated by the behavioural scenario. The costs from a long feeding cessation (t_d) depend on the energy content in prey (E_p) and the feeding rate (r_f), whereas the costs of a long (t_f) or fast (U_f) flight response depend on locomotor costs (C_L) and the change in fluking frequency (Δf). The relative energetic cost (E^* , fourth column) of a behavioural scenario is defined as the ratio of the energetic cost to the daily basal metabolic rate. Since energy budgets increase with body size, E^* facilitates interspecific comparisons

were temporarily lifted from the water to deploy bio-loggers, so the first hour following deployment was excluded to remove deployment response behaviours (Rojano-Doñate et al., 2018; Wisniewska et al., 2018). Each deployment was divided into hourly bins, excluding the first fraction of an hour, to generate species-specific empirical distributions. For example, a 1.5-hr deployment would yield one hourly feeding rate, counting all feeding events from 0.5 to 1.5 hr after deployment. Excluding the first fractional hour was intended to reduce the influence of potential changes in behaviour associated with tag deployment. Although data are sparse, available evidence suggests cetaceans have extremely high feeding success rates (Wisniewska et al., 2016), so we assumed all feeding events resulted in prey capture.

Energy acquired per feeding event (E_p) was modelled as a log-normal variable to match the distribution of resources in patchy environments (Benoit-Bird et al., 2019; Pagel et al., 1991; Sims et al., 2008; Sugihara, 1980). For odontocetes, the lognormal was parameterized using prey size and energy density data from the literature (Goldbogen et al., 2019, Tables S1–S9). Specifically, the lognormal mean and variance of E_p for each species were estimated as the

mean and variance of the logged product of prey size (in kg) and prey energy density (in kJ/kg), weighted by diet composition. We used acoustic prey mapping spatially and temporally linked with feeding whales to measure biomass densities available to rorquals (Goldbogen et al., 2019). Biomass density was modelled as a lognormal distribution and the energy per feeding event was then calculated as the product of biomass density (kg/m³; Cade et al., 2021), engulfment capacity (m³) and prey energy density (kJ/kg). Engulfment capacity was estimated from a morphological model (Goldbogen et al., 2010; Kahane-Rapport and Goldbogen, 2018; Potvin et al., 2012). We used krill energy density values based on *Thysanoessa spinifera* krill (3,800 kJ/kg) for all rorqual species except *B. bonaerensis*, which feeds on *Euphausia superba* (4,575 kJ/kg; Chenoweth, 2018; Färber-Lorda et al., 2009). Although rorquals feed on a variety of prey, we focused on krill-feeding individuals for two reasons. First, krill is the most common diet across species (Kawamura, 1980; Nemoto, 1970). Second, fish-feeding rorquals feed at lower and more variable rates and longer datasets are needed to properly quantify patterns in undisturbed feeding behaviour (Cade et al., 2016, 2020).

2.2 | Increased energy expenditure

The increase in energy expenditure due to the flight response was modelled as the additional locomotor costs from swimming at an elevated speed.

$$\Delta P_e(U_f) = \Delta f(U_f) C_L m, \quad (3)$$

where:

Δf is the increase in stroke frequency from cruising to flight speed in stroke/hr, a function of swim speed.

C_L is the mass-specific locomotor cost of a stroke in kJ stroke⁻¹ kg⁻¹.

m is the mass of the animal in kg.

The mass-specific locomotor cost of a stroke for cetaceans at cruising speed is estimated to be (Williams et al., 2017):

$$C_L = 1.46 \times 10^{-3} + 5.0 \times 10^{-7} m. \quad (4)$$

This relationship was derived using data from odontocete species massing 42 kg to 2,700 kg, and is therefore a source of uncertainty for larger odontocetes and mysticetes. Williams et al. (2017) also derived a locomotor cost relationship for maximum aerobic performance; however, the relationship had low explanatory power and produced unrealistic results when extrapolated to rorqual body sizes. We estimated the increase in stroke frequency by assuming cetaceans cruise at 1.5 m/s and swim efficiently by maintaining a Strouhal number of 0.3 (Rohr & Fish, 2004; Sato et al., 2007). The Strouhal number is a dimensionless ratio relating swimming speed to stroke amplitude and frequency ($St = \frac{Af}{U}$). Assuming a stroke amplitude of 1/5 body length (Bainbridge, 1958; Fish, 1998; Gough et al., 2019), a Strouhal number of 0.3 can be used to estimate stroke frequency as:

$$f = \frac{1.5U}{L} \times 3600, \quad (5)$$

where:

St is the Strouhal number (unitless).

A is the stroke amplitude in m.

f is the stroke frequency in strokes/s.

U is the swimming speed in m/s.

L is the length of the animal in m.

Substituting a cruising speed of 1.5 m/s (Gough et al., 2019; Sato et al., 2007; Watanabe et al., 2011) into Equation (5) gives the increase in stroke frequency (in strokes/hr) as:

$$\Delta f(U_f) = \frac{5400}{L} (U_f - 1.5). \quad (6)$$

There is considerable uncertainty associated with the values of many of the parameters in Equations 4 and 6. The slope of the scaling relationship for C_L (Equation 4) was not significant in the original study, likely due to small sample sizes, and the estimate for the change in fluking frequency depends on assumptions of a fixed cruising speed and Strouhal number (Equation 6). To account for this uncertainty, Equations 4 and 6 were treated as best estimates and C_L and Δf were drawn from gamma distributions with means equal to the best estimates. See the next section, *Sensitivity analysis*, for more details.

Combining Equations (1–3 and 6) yields the final, expanded model:

$$E = r_f E_p t_d + \frac{5400}{L} (U_f - 1.5) C_L m t_f. \quad (7)$$

2.3 | Sensitivity analysis

We conducted a sensitivity analysis using a global approach (i.e. one that samples all parameters simultaneously) to understand how altering one energetic parameter (r_f , E_p , Δf and C_L) changes the model result (E) across the full range of the other parameters. We generated a matrix of input values using Latin hypercube sampling (which produces random combinations of inputs evenly distributed throughout the parameter space), then calculated a vector of model outputs. A multiple linear regression of the outputs with respect to the inputs quantified the global sensitivity of the model to each parameter; that is, each coefficient describes the change in the output with respect to a change in one parameter after controlling for all other parameters. However, the four parameters considered in our sensitivity analysis are of greatly different magnitudes and in different units, so a standardization step is necessary for the regression coefficients to be comparable. We standardized the input matrix and the output vector by z-score, that is, the difference of the parameter value from the parameter mean divided by the standard deviation. Conducting the sensitivity analysis in this way allowed us to, for example, understand how changes in locomotor costs (C_L) affect model output (E) across all combinations of feeding rates (r_f), energy obtained per feeding event (E_p) and change in fluking frequency (Δf). This global approach is opposed to local sensitivity analyses where all other parameters are held at a fixed value (typically the mean) and

each parameter is varied individually, which describes sensitivity in a more limited context (Marino et al., 2008).

The distribution of hourly feeding rates (r_f) for each species was generated empirically by counting the number of feeding events (lunges in rorquals and buzzes in toothed whales) in each hour of each deployment, discarding the leading incomplete hour (see *Lost energy acquisition* above). The mean and standard deviation of the rorqual E_p distributions were based on prey mapping data from two California locations in the years 2011–2018 (Goldbogen et al., 2019). For odontocete E_p distributions, we fit a lognormal distribution to available prey data in the literature (Goldbogen et al., 2019). Both Δf and C_L were treated as gamma distributions with a mean equal to the parameter's best estimate (Equations 4 and 5). We sampled parameter values using Latin hypercube sampling with the pse R package (Chalom & de Prado 2017).

The model's sensitivity to energetic parameters is conditional on the behavioural response, for example, E will be more sensitive to r_f and E_p when t_d is very long relative to t_f . Therefore, we assessed model sensitivity for two different behavioural response scenarios that emphasize energy acquisition (i.e. greater t_d , lesser t_f and U_f) and energy expenditure (i.e. greater t_f and U_f , lesser t_d ; Table 3). The magnitude of the behavioural parameters in these scenarios was constrained to responses observed in controlled exposure experiments, and though they are not representative of the full spectrum of potential responses they provide a basis for comparing the sensitivity of energetic costs to lost feeding opportunities versus elevated locomotor effort. For each behavioural scenario and species, we generated a matrix of input values via Latin hypercube sampling and calculated a vector of model results. Inputs and outputs were standardized by z-score and we fit a linear regression to calculate a standardized coefficient for each parameter and species [$z(E) \sim z(r_f) + z(E_p) + z(\Delta f) + z(C_L)$] using the lm function in base R (R Core Team, 2018). Each coefficient, then, represents the change in model output if the parameter was increased by one standard deviation, after controlling for the other parameters.

2.4 | Relative energetic costs

As energy budgets increase with body size, a size-corrected metric is necessary for interspecific comparisons of energetic costs for a clade covering three orders of magnitude of body size. We used our model to estimate the absolute energetic costs associated with three behavioural scenarios corresponding to observed

TABLE 3 Behavioural scenarios used to test model sensitivity to energetic parameters. t_d , duration of feeding cessation; t_f , duration of flight response; U_f , swimming speed of flight response

Scenario	t_d (hr)	t_f (hr)	U_f (m/s)
Energy acquisition	4.0	0.25	3.0
Energy expenditure	1.0	1.0	5.0

mild, strong and extreme responses to sonar (Table 4). These costs were divided by daily basal metabolic demands, as predicted by allometric relationships between size and basal metabolic rate ($BMR = 293.1 \times [\text{mass (kg)}]^{0.75}$; Kleiber, 1975). The resulting metric, E^* , represents the relative energetic costs of a behavioural response with respect to body size and is only applicable to interspecific comparisons; it is not a meaningful metric for comparing intraspecific relative energetic costs for individuals differing in size or metabolic rates. E^* calculations for additional metabolic scaling relationships are reported in Supporting Information Appendix 1 (see Figures S5–S7; Maresh, 2014; Nagy et al., 1999; White & Seymour, 2003). All computational details (code, session information) are also available in Supporting Information Appendix 1.

3 | RESULTS

3.1 | Lost energy acquisition

Baseline rate of energy acquisition (P_a) was calculated as the product of the hourly feeding rate (r_f) and energy per feeding event (E_p ; Equation 2). r_f was zero-inflated and we modelled E_p with a lognormal distribution; therefore, P_a was zero-inflated and right-skewed (Figure 2). Rorquals and harbour porpoises had the greatest mass-specific energy acquisition rates while beaked whales and sperm whales had the lowest (Table 5). Hourly feeding rates were highly variable. Mean r_f ranged from 2.44 (first and third quartile, Q1–Q3: 0–3, short-finned pilot whale) to 110 feeding events per hour (Q1–Q3 19–169, harbour porpoise; Figure 3a). The first quartile of feeding rates was 0 for all but two species: harbour porpoises and sperm whales. E_p increased with body size such that bulk filter-feeding rorquals consumed multiple orders of magnitude more energy per feeding event than odontocetes (Table 5). However, the Q1–Q3 range of mass-specific energy acquired per feeding event overlapped among delphinids and balaenopterids, with substantially lower values for ziphiids and physeterids (Figure 3b).

3.2 | Sensitivity analysis

Parameters associated with lost energy acquisition (r_f , E_p) were more influential than the energy expenditure parameters (Δf , C_L) across species groups in both behavioural scenarios. In the scenario

TABLE 4 Behavioural scenarios used to model relative energetic costs. t_d , duration of feeding cessation; t_f , duration of flight response; U_f , swimming speed of flight response

Scenario	t_d (hr)	t_f (hr)	U_f (m/s)
Mild response	1	0.25	2.5
Strong response	2	0.5	3.5
Extreme response	8	2	5




FIGURE 2 Example of baseline rate of energy acquisition (ΔE_a , main plot) for a blue whale as the product of lognormally distributed prey energy content (E_p) and empirically measured hourly feeding rates (r_f ; inset plots). Mean P_a indicated with dashed line. Similar distributions were calculated for all species in the study (see Supporting Information)

emphasizing feeding cessation ($t_d = 4$ hr, $t_f = 0.25$ hr, $U_f = 3.0$ m/s), sensitivity coefficients were greater for r_f and E_p than Δf and C_L with no overlap in the 95% confidence intervals (Figure 4a). Model sensitivity to energy expenditure parameters increased in the elevated flight scenario ($t_d = 1$ hr, $t_f = 1$ hr, $U_f = 5.0$ m/s), but did not fully exceed sensitivity to energy acquisition parameters (Figure 4b). In this scenario, E_p was the most sensitive parameter for most species followed by C_L and r_f . Rorquals were most sensitive to E_p (95% CI 0.524–0.593) followed by C_L (95% CI 0.389–0.456) and r_f (95% CI 0.165–0.220). Similarly, porpoises and dolphins were most sensitive to E_p (95% CI 0.499–0.575) followed by C_L (95% CI 0.175–0.247) and r_f (95% CI 0.065–0.144). Only the beaked and sperm whales were most sensitive to an energy expenditure parameter, C_L (95% CI 0.743–0.798), followed by E_p (95% CI 0.254–0.308) and Δf (95% CI 0.103–0.155).

3.3 | Relative energetic costs

Relative energetic costs (E^*) were concave upward with respect to body size, indicating that the smallest and largest species face greater short-term relative energetic costs (Figure 5). Within each scenario, the greatest median E^* values were associated with humpback and blue whales and the lowest with short-finned pilot whales

TABLE 5 Energy acquisition parameters (feeding rate in events/hr: r_f , energy acquired per prey capture event: E_p , baseline energy intake rate: P_a , and mass-specific rate: P_a/m) presented as mean and Q1–Q3. Species ordered by size, mysticetes in bold.

Species (n individuals, hours)	r_f (hr ⁻¹)	E_p (kJ)	P_a (kJ/hr)	P_a/m (kJ hr ⁻¹ kg ⁻¹)
<i>P. phocoena</i> (8, 161)	110 (19–169)	14.9 (8.20–27.2)	2.40×10^3 (186–2.86 $\times 10^3$)	74.5
<i>G. griseus</i> (11, 39)	11.7 (0–17.5)	282.0 (215.0–369.0)	3.26×10^3 (0.0–4.31 $\times 10^3$)	8.57
<i>M. densirostris</i> (14, 105)	12.8 (0–23)	218.0 (152.0–313.0)	2.97×10^3 (0.0–4.81 $\times 10^3$)	3.95
<i>G. macrorhynchus</i> (2, 9)	2.44 (0–3)	1.31×10^3 (732.0–2.33 $\times 10^3$)	3.34×10^3 (0.0–2.74 $\times 10^3$)	3.80
<i>G. melas</i> (9, 101)	7.37 (0–13)	1.31×10^3 (732.0–2.33 $\times 10^3$)	1.25×10^4 (0.0–1.52 $\times 10^4$)	11.1
<i>Z. cavirostris</i> (3, 53)	10.8 (0–21)	419.0 (223.0–786.0)	6.89×10^3 (0.0–7.31 $\times 10^3$)	2.41
<i>B. bonaerensis</i> (6, 121)	48.4 (0–62)	2.82×10^3 (2.28 $\times 10^3$ –3.50 $\times 10^3$)	1.53×10^5 (2.25 $\times 10^3$ –1.97 $\times 10^5$)	20.6
<i>P. macrocephalus</i> (29, 237)	11.4 (4–17)	2.70×10^3 (1.57 $\times 10^3$ –4.64 $\times 10^3$)	4.19×10^4 (8.94 $\times 10^3$ –5.61 $\times 10^4$)	2.83
<i>M. novaeangliae</i> (28, 229)	17.0 (0–26)	9.29×10^4 (7.42 $\times 10^4$ –1.16 $\times 10^5$)	1.68×10^6 (0.0–2.68 $\times 10^6$)	45.4
<i>B. physalus</i> (4, 39)	14.8 (0–26)	1.44×10^5 (1.16 $\times 10^5$ –1.78 $\times 10^5$)	2.07×10^6 (0.0–3.63 $\times 10^6$)	42.0
<i>B. musculus</i> (21, 243)	14.3 (0–24)	3.13×10^5 (2.46 $\times 10^5$ –3.98 $\times 10^5$)	4.98×10^6 (0.0–7.86 $\times 10^6$)	51.4




FIGURE 3 (A) hourly feeding rates (r_f , mean and Q1–Q3) were greatest for the smallest odontocete (harbour porpoise) and mysticetes (minke whale). (b) Mass-specific energy acquired per feeding event (mass-specific E_p , mean and Q1–Q3) increased with body size with the exception of beaked and sperm whales. Families Phocoenidae and Delphinidae in blue triangles, Ziphiidae and Physeteridae in orange squares, and Balaenopteridae in green circles. Species ordered by size; see Table 2 for abbreviations




FIGURE 4 Sensitivity analyses indicated parameters associated with energy acquisition (energy from prey, E_p ; feeding rate, r_f) were more influential than energy expenditure parameters (locomotor cost, C_L ; change in stroke frequency, Δf) in behavioural scenarios emphasizing both energy acquisition (a) and energy expenditure (b; Table 3). Families Phocoenidae and Delphinidae in blue, Ziphiidae and Physeteridae in orange, and Balaenopteridae in green. See text for details




FIGURE 5 The relative energetic costs (E^*) of three behavioural responses (Table 4) with respect to body size (points and lines: median and Q1–Q3). The concave pattern indicates very large and small cetaceans face greater immediate energetic costs relative to their size for a given behavioural response than intermediate-sized species. The relative energetic costs of a mild response by large rorquals exceeded those of an extreme response by short-finned pilot whales and beaked whales. Families Phocoenidae and Delphinidae in blue, Ziphiidae and Physeteridae in orange, and Balaenopteridae in green

and the beaked whales. Across behavioural scenarios, the relative energetic cost of a mild behavioural response by a blue whale ($t_d = 1$ hr, $t_f = 0.25$ hr, $U_f = 2.5$ m/s) was significantly greater than an extreme behavioural response ($t_d = 8$ hr, $t_f = 2.0$ hr, $U_f = 5.0$ m/s) by a Cuvier's beaked whale (t test, $p < 0.001$).

4 | DISCUSSION

We modelled the energetic costs of sonar disturbance to cetaceans using measurements of undisturbed feeding rates, empirical prey characteristics and energetic costs from changes in locomotion based on metabolic allometry and biomechanical constraints. Lost feeding opportunities generally had a greater impact on energetic costs than elevated locomotion. The importance of feeding cessation relative to increased locomotor costs during disturbances was previously identified for minke and killer whales (Christiansen et al., 2013; Noren et al., 2016; Williams et al., 2006). This work generalizes those findings, as energetic costs were more sensitive

to lost feeding opportunities for most species and behavioural response scenarios. The only exceptions were the two beaked whales and the sperm whale in a scenario emphasizing elevated locomotion. Considering the model's sensitivity to locomotor costs for those species and the magnitude of flight responses observed in Cuvier's beaked whale during controlled exposure experiments (DeRuiter et al., 2013), cost of transport for large, deep-diving odontocetes is a critical data gap. The harbour porpoise and orqualws faced the greatest immediate relative energetic costs (E^*) due to their high energy intake potential. As such, the relative energetic costs incurred by a blue whale exhibiting a 'mild' behavioural response exceed those of an 'extreme' response by a Cuvier's beaked whale (Figure 5).

There is an apparent correlation between species' relative energetic costs from disturbance (E^*) and their life history, behaviour and physiology (i.e. their 'pace of life'). The pace of life syndrome concept describes a slow-fast continuum based on covarying biological traits (Réale et al., 2010). Traits of fast-paced species include short life spans, high reproductive output and lower sociability. Concordantly, the high E^* species, porpoises and orqualws, have reduced longevity




FIGURE 6 (a) The energetic models presented here fill a gap in the PCoD framework along the behavioural-energetic pathway. Earlier behavioural response studies quantified the relationships, and uncertainty, between the exposure to a stressor (sonar) and the subsequent behaviour changes (feeding cessation and elevated locomotion). The present study predicts the energetic costs of these responses, connecting behavioural changes to health. Future empirical work is needed to comprehensively understand how reduced energy stores affect vital rates. (b) Species with the greatest relative energetic costs from disturbance (porpoises, orqualws) have low longevity for their size while those with lower relative energetic costs (larger odontocetes, such as pilot whales and sperm whales) have greater longevity for their size. The y-axis shows the ratio of female cetaceans' longevity (Trites & Pauly, 1998) to allometrically predicted longevity (Speakman, 2005). Species included in the present study are labelled; see table 2 in Trites and Pauly (1998) for complete list of species. Families Phocoenidae and Delphinidae in blue, Physeteridae in orange, and Balaenopteridae in green. (c) In addition to longevity, orqualws and porpoises exhibit fast pace of life traits such as rapid reproduction and high feeding effort. Larger odontocetes, such as beaked and sperm whales, reproduce more slowly and exhibit greater sociality, typical of a slow pace of life. Population sizes of species with a fast pace of life are often more sensitive to reduced reproductive success, whereas survival is a greater concern for those with a slow pace of life. Illustrations by Alex Boersma

for their size (Figure 6b) and rapid reproductive cycles (Figure 6c; Speakman, 2005; Trites & Pauly, 1998). Conversely, low E^* species, such as delphinids and sperm whales, form complex social networks and their parental care extends for years longer than rorquals (Lockyer, 2007; May-Collado et al., 2007). Overall, species with high relative energetic costs from disturbance have a fast pace of life and vice versa. This has conservation implications because pace-of-life correlates with population dynamics. Specifically, population growth of fast-paced species is more sensitive to changes in fecundity, whereas slow-paced species are more sensitive to changes in survival (Oli, 2004; Stahl & Oli, 2006).

Our model can be applied in the Population Consequences of Disturbance (PCoD) framework to predict the immediate energetic costs for behavioural responses. Table 6 illustrates how resource managers could use the model for a variety of species and contexts. These energetic costs impact individuals' energy stores and body condition if they cannot behaviourally or physiologically compensate. Future research specific to species' foraging ecology and life history is therefore critically needed (Booth, 2020; Pirodda et al., 2021). For example, blue whales face enormous immediate energetic costs from disturbance due to their remarkable foraging efficiency. However, that efficiency could also facilitate rapid compensation, depending on their time budgets (can they make up feeding later?) and the distribution of their prey (can they find another prey patch?). These unresolved questions may be answered by longer duration, high-resolution tags (Calambokidis et al., 2019, 2020) and prey mapping at spatial scales relevant to foraging whales (Cade et al., 2021). As a species with a fast pace of life, their population growth is likely more sensitive to changes in reproduction than survival. Therefore, research on the transfer function linking energy stores to reproductive success should be prioritized.

Our model was designed to be broadly applicable to other species and sources of disturbance. Cetaceans' behavioural responses to disturbance are highly variable and context-dependent (Ellison et al., 2012). By parameterizing the behavioural response as time

displaced from feeding (t_d), time fleeing the sound source (t_f), and swimming speed (U_f), resource managers can draw on expertise in their system to predict short-term energetic costs from disturbance, regardless of the sound source. Our model is equally applicable to whale-watching (Senigaglia et al., 2016) and underwater pile driving (Thompson et al., 2010) as it is to sonar, allowing resource managers to understand the energetic costs of anticipated behavioural responses. Furthermore, our scaling approach facilitates assessments of understudied species for which empirical energetic data are unavailable. For example, PCoD models for Cook Inlet belugas *Delphinapterus leucas* face data deficiencies that necessitate expert solicitation to estimate the magnitude of energetic costs of disturbance (Tollit et al., 2016). Our model predictions for similarly sized species complement expert opinion and serve as an additional input for conservation planning.

Here, we have added to the PCoD framework by connecting behaviour to health using empirical foraging data. Specifically, we build on the behavioural-energetic pathway. Other pathways may be triggered by direct physiological responses and/or affect other aspects of health than body condition, such as stress levels, immune status and organ condition (Schwacke et al., 2014). Along the behavioural-energetic pathway, earlier behavioural response studies used controlled field experiments to evaluate altered foraging and movement behaviours (Table 1). Downstream, further research is needed to quantify the connections between health (i.e. body condition) and vital rates, most critically survival and fecundity (Christiansen & Lusseau, 2015; Christiansen et al., 2018). Our model applies an extensive dataset to address the gap between foraging behaviour and vital rates. In the absence of these data, previous work was limited by unknown energy intake rates in undisturbed conditions. Although the capacity to compensate for these energetic costs remains unknown, we uncover a correlation between the magnitude of short-term (minutes to hours) energetic costs with species' pace of life. At time-scales relevant to population dynamics (months to years), the chronic energetic effects of disturbance depend not only on

TABLE 6 Example applications of the present model to humpback whales in Hawaii and blue whales and Cuvier's beaked whale in California. Behaviour parameters (first column) may be drawn from behavioural response studies when available, otherwise from expert opinion. The 0 hr displacement from feeding (t_d) for the Hawaii humpback whale reflects its capital breeding strategy and the behaviour parameters for the blue whale (Southall et al., 2019) and Cuvier's beaked whale (DeRuiter et al., 2013) are based on behavioural response studies. The immediate energy costs (second column) establish the magnitude of compensation necessary to avoid body condition deterioration. Options for compensation (third column) depend on foraging ecology and reproductive strategy. The species' pace of life (fast for humpback and blue whales, slow for Cuvier's beaked whale) suggests the most sensitive vital rate (fourth column) for consideration in full PCoD models

	Behaviour parameters	Energy costs (kJ)	Compensation options	Key vital rate
Humpback whale (HI)	$t_d = 0$ hr $t_f = 0.3$ hr $U_f = 2.8$ m/s	8.17×10^4 ($5.07 \times 10^4 - 1.40 \times 10^5$)	Terminate lactation	Calf survival
Blue whale (CA)	$t_d = 0.25$ hr $t_f = 0.2$ hr $U_f = 3.0$ m/s	1.25×10^6 ($4.20 \times 10^5 - 2.01 \times 10^6$)	Additional feeding, delay migration	Gestation termination
Cuvier's beaked whale (CA)	$t_d = 7.6$ hr $t_f = 1.7$ hr $U_f = 3.1$ m/s	3.30×10^4 ($1.48 \times 10^4 - 6.69 \times 10^4$)	Additional feeding, terminate gestation/lactation	Adult survival

the frequency and magnitude of these costs but also on individuals' ability to compensate. Fast-paced species (porpoises, rorquals) face relatively greater short-term costs, but whether that translates to long-term deficits remains unknown and warrants future work. Using empirical data to quantify short-term energetic costs allows improved prediction of the tipping points where chronic disturbance leads to population declines.

ACKNOWLEDGEMENTS

The study was funded in part by the U.S. Office of Naval Research (grant N00014-16-1-2477). Support for M.F.C. was provided by Stanford University's William R. and Sara Hart Kimball Fellowship and a Data Science Scholar Fellowship. We thank Elliott Hazen for guidance on the sensitivity analysis. We are grateful to the associate editor and four reviewers for their constructive critiques, which greatly improved the manuscript.

AUTHORS' CONTRIBUTIONS

M.F.C., M.S.S. and J.A.G. conceived the study; All authors contributed to study design and provided critical feedback on the manuscript; M.F.C. was responsible for writing the manuscript and data analyses.

DATA AVAILABILITY STATEMENT

Data available via the Dryad Digital Repository <https://doi.org/10.5061/dryad.pvmcvdnq> (Czapanskiy et al., 2021a). An R Markdown file with all computational details for running the analysis and producing the figures and tables is archived on Zenodo <https://doi.org/10.5281/zenodo.4646110> (Czapanskiy et al., 2021b).

ORCID

Max F. Czapanskiy  <https://orcid.org/0000-0002-6302-905X>
 Matthew S. Savoca  <https://orcid.org/0000-0002-7318-4977>
 William T. Gough  <https://orcid.org/0000-0003-2701-5299>
 Paolo S. Segre  <https://orcid.org/0000-0002-2396-2670>
 Danuta M. Wisniewska  <https://orcid.org/0000-0002-3599-7440>
 David E. Cade  <https://orcid.org/0000-0003-3641-1242>
 Jeremy A. Goldbogen  <https://orcid.org/0000-0002-4170-7294>

REFERENCES

- Antunes, R., Kvadsheim, P. H., Lam, F. P. A., Tyack, P. L., Thomas, L., Wensveen, P. J., & Miller, P. J. O. (2014). High Thresholds for avoidance of sonar by free-ranging long-finned pilot whales (*Globicephala melas*). *Marine Pollution Bulletin*, 83(1), 165–180. <https://doi.org/10.1016/j.marpolbul.2014.03.056>
- Bainbridge, R. (1958). The speed of swimming of fish as related to size and to the frequency and amplitude of the tail beat. *Journal of Experimental Biology*, 35(1), 109–133. <https://doi.org/10.1242/jeb.35.1.109>
- Barber, J. R., Crooks, K. R., & Frstrup, K. M. (2010). The costs of chronic noise exposure for terrestrial organisms. *Trends in Ecology & Evolution*, 25(3), 180–189. <https://doi.org/10.1016/j.tree.2009.08.002>
- Benoit-Bird, K. J., & Lawson, G. L. (2016). Ecological insights from pelagic habitats acquired using active acoustic techniques. *Annual Review of Marine Science*, 8(1), 463–490. <https://doi.org/10.1146/annurev-marine-122414-034001>
- Benoit-Bird, K. J., Waluk, C. M., & Ryan, J. P. (2019). Forage species swarm in response to coastal upwelling. *Geophysical Research Letters*, 46(3), 1537–1546. <https://doi.org/10.1029/2018GL081603>
- Blix, A. S., & Folkow, L. P. (1995). Daily energy expenditure in free living minke whales. *Acta Physiologica Scandinavica*, 153(1), 61–66. <https://doi.org/10.1111/j.1748-1716.1995.tb09834.x>
- Booth, C. G. (2020). Food for thought: Harbor porpoise foraging behavior and diet inform vulnerability to disturbance. *Marine Mammal Science*, 36(1), 195–208. <https://doi.org/10.1111/mms.12632>
- Cade, D. E., Carey, N., Domenici, P., Potvin, J., & Goldbogen, J. A. (2020). Predator-informed looming stimulus experiments reveal how large filter feeding whales capture highly maneuverable forage fish. *Proceedings of the National Academy of Sciences of the United States of America*, 117(1), 472–478. <https://doi.org/10.1073/pnas.1911099116>
- Cade, D. E., Friedlaender, A. S., Calambokidis, J., & Goldbogen, J. A. (2016). Kinematic diversity in Rorqual whale feeding mechanisms. *Current Biology*, 26(19), 2617–2624. <https://doi.org/10.1016/j.cub.2016.07.037>
- Cade, D. E., Seakamela, S. M., Findlay, K. P., Fukunaga, J., Kahane-Rapport, S. R., Warren, J. D., Calambokidis, J., Fahlbusch, J. A., Friedlaender, A. S., Hazen, E. L., Kotze, D., McCue, S., Meyer, M., Oestreich, W. K., Oudejans, M. G., Wilke, C., & Goldbogen, J. A. (2021). Predator-scale spatial analysis of intra-patch prey distribution reveals the energetic drivers of Rorqual whale super-group formation. *Functional Ecology*, 35(4), 894–908. <https://doi.org/10.1111/1365-2435.13763>
- Calambokidis, J., Fahlbusch, J. A., Szesciorka, A. R., Sirovic, A., Southall, B. L., Cade, D. E., Friedlaender, A. S., Oestreich, W., & Goldbogen, J. A. (2020). *New insights into blue, fin, and humpback whale feeding, movements, and impacts of human activities from medium-duration archival tags* (Conference presentation). Ocean Sciences Meeting 2020.
- Calambokidis, J., Fahlbusch, J. A., Szesciorka, A. R., Southall, B. L., Cade, D. E., Friedlaender, A. S., & Goldbogen, J. A. (2019). Differential vulnerability to ship strikes between day and night for blue, fin, and humpback whales based on dive and movement data from medium duration archival tags. *Frontiers in Marine Science*, 6. <https://doi.org/10.3389/fmars.2019.00543>
- Chalom, A., & de Prado, P. I. K. L. (2017). *Pse: Parameter Space Exploration with Latin Hypercubes*. R package version 0.4.7. Retrieved from <https://cran.r-project.org/package=pse>
- Chenoweth, E. M. (2018). *Bioenergetic and economic impacts of humpback whale depredation at salmon hatchery release sites* (Doctoral dissertation). University of Alaska Fairbanks.
- Christiansen, F., & Lusseau, D. (2015). Linking behavior to vital rates to measure the effects of non-lethal disturbance on wildlife. *Conservation Letters*, 8(6), 424–431. <https://doi.org/10.1111/conl.12166>
- Christiansen, F., Rasmussen, M. H., & Lusseau, D. (2013). Inferring activity budgets in wild animals to estimate the consequences of disturbances. *Behavioral Ecology*, 24(6), 1415–1425. <https://doi.org/10.1093/beheco/art086>
- Christiansen, F., Vivier, F., Charlton, C., Ward, R., Amerson, A., Burnell, S., & Bejder, L. (2018). Maternal body size and condition determine calf growth rates in Southern Right Whales. *Marine Ecology Progress Series*, 592, 267–281. <https://doi.org/10.3354/meps12522>
- Clarke, M. R. (1996). Cephalopods as Prey. III. Cetaceans. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 351(1343), 1053–1065. <https://doi.org/10.1098/rstb.1996.0093>
- Cox, T. M., Ragen, T. J., Read, A. J., Vos, E., Baird, R. W., Balcomb, K., Barlow, J., Caldwell, J., Cranford, T., Crum, L., Angela D'Amico, G., D'Spain, A., Fernández, J., Finneran, R., Gentry, W., Gerth, F. G., Hildebrand, J. A., Dorian Houser, T., Hullah, P. D., ... Benner, L. (2006). Understanding the Impacts of Anthropogenic Sound on Beaked Whales. *Journal of Cetacean Research and Management*, 7(3), 177–187.

- Czapanskiy, M. F., Savoca, M. S., Gough, W. T., Segre, P. S., Wisniewska, D. M., Cade, D. E., & Goldbogen, J. A. (2021a). Data from: Modelling short-term energetic costs of sonar disturbance to cetaceans using high-resolution foraging data. *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.pvmcndnkq>
- Czapanskiy, M. F., Savoca, M. S., Gough, W. T., Segre, P. S., Wisniewska, D. M., Cade, D. E., & Goldbogen, J. A. (2021b). Supplemental Information for: "Modelling short-term energetic costs of sonar disturbance to cetaceans using high-resolution foraging data". *Zenodo*, <https://doi.org/10.5281/zenodo.4646109>
- D'Amico, A., Gisiner, R. C., Ketten, D. R., Hammock, J. A., Johnson, C., Tyack, P. L., & Mead, J. (2009). Beaked whale strandings and naval exercises. *Aquatic Mammals*, *35*(4), 452–472. <https://doi.org/10.1578/AM.35.4.2009.452>
- DeRuiter, S. L., Southall, B. L., Calambokidis, J., Zimmer, W. M. X., Sadykova, D., Falcone, E. A., Friedlaender, A. S., Joseph, J. E., Moretti, D., Schorr, G. S., Thomas, L., & Tyack, P. L. (2013). First direct measurements of behavioural responses by Cuvier's beaked whales to mid-frequency active sonar. *Biology Letters*, *9*(4), 20130223. <https://doi.org/10.1098/rsbl.2013.0223>
- Dial, K. P., Greene, E., & Irschick, D. J. (2008). Allometry of behavior. *Trends in Ecology & Evolution*, *23*(7), 394–401. <https://doi.org/10.1016/j.tree.2008.03.005>
- Duarte, C. M., Chapuis, L., Collin, S. P., Costa, D. P., Devassy, R. P., Eguiluz, V. M., Erbe, C., Gordon, T. A. C., Halpern, B. S., Harding, H. R., Havlik, M. N., Meekan, M., Merchant, N. D., Miksis-Olds, J. L., Parsons, M., Predragovic, M., Radford, A. N., Radford, C. A., Simpson, S. D., ... Juanes, F. (2021). The soundscape of the Anthropocene ocean. *Science*, *371*(6529), eaba4658. <https://doi.org/10.1126/science.aba4658>
- Ellison, W. T., Southall, B. L., Clark, C. W., & Frankel, A. S. (2012). A new context-based approach to assess marine mammal behavioral responses to anthropogenic sounds. *Conservation Biology*, *26*(1), 21–28. <https://doi.org/10.1111/j.1523-1739.2011.01803.x>
- Fahlman, A., van der Hoop, J., Moore, M. J., Levine, G., Rocho-Levine, J., & Brodsky, M. (2016). Estimating energetics in cetaceans from respiratory frequency: Why we need to understand physiology. *Biology Open*, *5*(4), 436–442. <https://doi.org/10.1242/bio.017251>
- Färber-Lorda, J., Gaudy, R., & Mayzaud, P. (2009). Elemental composition, biochemical composition and caloric value of Antarctic krill: Implications in energetics and carbon balances. *Journal of Marine Systems*, *78*(4), 518–524. <https://doi.org/10.1016/j.jmarsys.2008.12.021>
- Farmer, N. A., Noren, D. P., Fougères, E. M., Machernis, A., & Baker, K. (2018). Resilience of the endangered sperm whale *Physeter macrocephalus* to foraging disturbance in the gulf of Mexico, USA: A bio-energetic approach. *Marine Ecology Progress Series*, *589*, 241–261. <https://doi.org/10.3354/meps12457>
- Fish, F. E. (1998). Comparative kinematics and hydrodynamics of Odontocete cetaceans: Morphological and ecological correlates with swimming performance. *Journal of Experimental Biology*, *201*(20), 2867–2877. <https://doi.org/10.1242/jeb.201.20.2867>
- Frid, A., & Dill, L. M. (2002). Human-caused disturbance stimuli as a form of predation risk. *Conservation Ecology*, *6*(1), 11–26. <https://doi.org/10.5751/ES-00404-060111>
- Friedlaender, A. S., Hazen, E. L., Goldbogen, J. A., Stimpert, A. K., Calambokidis, J., & Southall, B. L. (2016). Prey-mediated behavioral responses of feeding blue whales in controlled sound exposure experiments. *Ecological Applications*, *26*(4), 1075–1085. <https://doi.org/10.1002/15-0783>
- Frisk, G. V. (2012). Noiseonomics: The relationship between ambient noise levels in the sea and global economic trends. *Scientific Reports*, *2*, 437. <https://doi.org/10.1038/srep00437>
- Gill, J. A., Norris, K., & Sutherland, W. J. (2001). Why behavioural responses may not reflect the population consequences of human disturbance. *Biological Conservation*, *97*(2), 265–268. [https://doi.org/10.1016/S0006-3207\(00\)00002-1](https://doi.org/10.1016/S0006-3207(00)00002-1)
- Goldbogen, J. A., Cade, D. E., Calambokidis, J., Friedlaender, A. S., Potvin, J., Segre, P. S., & Werth, A. J. (2017). How baleen whales feed: The biomechanics of engulfment and filtration. *Annual Review of Marine Science*, *9*(1), 367–386. <https://doi.org/10.1146/annurev-marine-122414-033905>
- Goldbogen, J. A., Cade, D. E., Wisniewska, D. M., Potvin, J., Segre, P. S., Savoca, M. S., Hazen, E. L., Czapanskiy, M. F., Kahane-Rapport, S. R., DeRuiter, S. L., Gero, S., Tønnesen, P., Gough, W. T., Hanson, M. B., Holt, M. M., Jensen, F. H., Simon, M., Stimpert, A. K., Arranz, P., ... Pyenson, N. D. (2019). Why whales are big but not bigger: Physiological drivers and ecological limits in the age of ocean giants. *Science*, *366*(6471), 1367–1372. <https://doi.org/10.1126/science.aax9044>
- Goldbogen, J. A., Potvin, J., & Shadwick, R. E. (2010). Skull and buccal cavity allometry increase mass-specific engulfment capacity in fin whales. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1683), 861–868. <https://doi.org/10.1098/rspb.2009.1680>
- Goldbogen, J. A., Southall, B. L., DeRuiter, S. L., Calambokidis, J., Friedlaender, A. S., Hazen, E. L., Falcone, E. A., Schorr, G. S., Douglas, A., Moretti, D. J., Kyburg, C., McKenna, M. F., & Tyack, P. L. (2013). Blue whales respond to simulated mid-frequency military sonar. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1765), 20130657. <https://doi.org/10.1098/rspb.2013.0657>
- Gough, W. T., Segre, P. S., Bierlich, K. C., Cade, D. E., Potvin, J., Fish, F. E., Dale, J., di Clemente, J., Friedlaender, A. S., Johnston, D. W., Kahane-Rapport, S. R., Kennedy, J., Long, J. H., Oudejans, M., Penry, G., Savoca, M. S., Simon, M., Videsen, S. K. A., Visser, F., ... Goldbogen, J. A. (2019). Scaling of swimming performance in baleen whales. *Journal of Experimental Biology*, *222*(20). <https://doi.org/10.1242/jeb.204172>
- Griffin, S. C., Valois, T., Taper, M. L., & Scott Mills, L. (2007). Effects of tourists on behavior and demography of olympic marmots. *Conservation Biology*, *21*(4), 1070–1081. <https://doi.org/10.1111/j.1523-1739.2007.00688.x>
- Harris, C. M., Thomas, L., Falcone, E. A., Hildebrand, J., Houser, D., Kvadsheim, P. H., Lam, F.-P., Miller, P. J. O., Moretti, D. J., Read, A. J., Slabbekoorn, H., Southall, B. L., Tyack, P. L., Wartzok, D., & Janik, V. M. (2017). Marine mammals and sonar: Dose-response studies, the risk-disturbance hypothesis and the role of exposure Context. *Journal of Ecology*, *55*(1), 396–404.
- Harwood, J., King, S., Booth, C., Donovan, C., Schick, R. S., Thomas, L., & New, L. (2016). Understanding the population consequences of acoustic disturbance for marine mammals. In A. N. Popper & A. Hawkins (Eds.), *The effects of noise on aquatic life II* (pp. 417–423). Springer New York.
- Hildebrand, J. A. (2009). Anthropogenic and natural sources of ambient noise in the ocean. *Marine Ecology Progress Series*, *395*, 5–20. <https://doi.org/10.3354/meps08353>
- Hin, V., Harwood, J., & de Roos, A. M. (2019). Bio-energetic modeling of medium-sized cetaceans shows high sensitivity to disturbance in seasons of low resource supply. *Ecological Applications*, *29*(5), e01903. <https://doi.org/10.1002/eap.1903>
- van der Hoop, J. M., Nousek-McGregor, A. E., Nowacek, D. P., Parks, S. E., Tyack, P., & Madsen, P. T. (2019). Foraging rates of ram-filtering north Atlantic right whales. *Functional Ecology*, *33*(7), 1290–1306. <https://doi.org/10.1111/1365-2435.13357>
- Injaia, A. S., Poon, L. Y., & Patricelli, G. L. (2018). Effects of experimental anthropogenic noise on avian settlement patterns and reproductive success. *Behavioral Ecology*, *29*(5), 1181–1189. <https://doi.org/10.1093/beheco/ary097>
- Jensen, F. H., Beedholm, K., Wahlberg, M., Bejder, L., & Madsen, P. T. (2012). Estimated communication range and energetic cost of bottlenose dolphin whistles in a tropical habitat. *The Journal of*

- the *Acoustical Society of America*, 131(1), 582–592. <https://doi.org/10.1121/1.3662067>
- Jepson, P. D., Arbelo, M., Deaville, R., Patterson, I. A. P., Castro, P., Baker, J. R., Degollada, E., Ross, H. M., Herráez, P., Pocknell, A. M., Rodríguez, F., Howie, F. E., Espinosa, A., Reid, R. J., Jaber, J. R., Martin, V., Cunningham, A. A., & Fernández, A. (2003). Gas-bubble lesions in stranded cetaceans. *Nature*, 425(6958), 575–576. <https://doi.org/10.1038/425575a>
- Johnson, M., Madsen, P. T., Zimmer, W. M. X., Aguilar, N., Aguilar de Soto, N., & Tyack, P. L. (2004). Beaked whales echolocate on prey. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271, S383–S386. <https://doi.org/10.1098/rsbl.2004.0208>
- Kahane-Rappoport, S. R., & Goldbogen, J. A. (2018). Allometric scaling of morphology and engulfment capacity in Rorqual Whales. *Journal of Morphology*, 279(9), 1256–1268. <https://doi.org/10.1002/jmor.20846>
- Kaplan, M. B., & Solomon, S. (2016). A coming boom in commercial shipping? The potential for rapid growth of noise from commercial ships by 2030. *Marine Policy*, 73, 119–121. <https://doi.org/10.1016/j.marpol.2016.07.024>
- Kawamura, A. (1980). A review of food of balaeopterid whales. *The Scientific Reports of the Whales Research Institute*, 32, 155–197.
- King, S. L., Schick, R. S., Donovan, C., Booth, C. G., Burgman, M., Thomas, L., & Harwood, J. (2016). An interim framework for assessing the population consequences of disturbance. *Methods in Ecology and Evolution*, 6(10), 1150–1158. <https://doi.org/10.1111/2041-210X.12411>
- Kleiber, M. (1975). *The fire of life: An introduction to animal energetics*. R. E. Krieger.
- Kleist, N. J., Guralnick, R. P., Cruz, A., Lowry, C. A., & Francis, C. D. (2018). Chronic anthropogenic noise disrupts glucocorticoid signaling and has multiple effects on fitness in an avian community. *Proceedings of the National Academy of Sciences of the United States of America*, 115(4), E648–E657. <https://doi.org/10.1073/pnas.1709200115>
- Kvadsheim, P. H., DeRuiter, S., Sivle, L. D., Goldbogen, J. A., Roland-Hansen, R., Miller, P. J. O., Lam, F.-P., Calambokidis, J., Friedlaender, A., Visser, F., Tyack, P. L., Kleivane, L., & Southall, B. (2017). Avoidance responses of minke whales to 1–4 KHz naval sonar. *Marine Pollution Bulletin*, 121(1–2), 60–68. <https://doi.org/10.1016/j.marpol.2017.05.037>
- Lindberg, D. R., & Pyenson, N. D. (2007). Things that go bump in the night: Evolutionary interactions between cephalopods and cetaceans in the tertiary. *Lethaia*, 40(4), 335–343. <https://doi.org/10.1111/j.1502-3931.2007.00032.x>
- Lockyer, C. (2007). All creatures great and smaller: A study in cetacean life history energetics. *Journal of the Marine Biological Association of the United Kingdom*, 87(4), 1035–1045. <https://doi.org/10.1017/S0025315407054720>
- Maresh, J. L. (2014). *Bioenergetics of marine mammals: The influence of body size, reproductive status, locomotion and phylogeny on metabolism* (Doctoral dissertation). University of California.
- Marino, S., Hogue, I. B., Ray, C. J., & Kirschner, D. E. (2008). A methodology for performing global uncertainty and sensitivity analysis in systems biology. *Journal of Theoretical Biology*, 254(1), 178–196. <https://doi.org/10.1016/j.jtbi.2008.04.011>
- May-Collado, L. J., Agnarsson, I., & Wartzok, D. (2007). Phylogenetic review of tonal sound production in whales in relation to sociality. *BMC Evolutionary Biology*, 7(1), 136. <https://doi.org/10.1186/1471-2148-7-136>
- McDonald, M. A., Hildebrand, J. A., & Wiggins, S. M. (2006). Increases in deep ocean ambient noise in the northeast Pacific West of San Nicolas Island, California. *The Journal of the Acoustical Society of America*, 120(2), 711–718. <https://doi.org/10.1121/1.2216565>
- Miller, P. J. O., Antunes, R. N., Wensveen, P. J., Samarra, F. I. P., Alves, A. C., Tyack, P. L., Kvadsheim, P. H., Kleivane, L., Lam, F.-P., Ainslie, M. A., & Thomas, L. (2014). Dose-response relationships for the onset of avoidance of sonar by free-ranging killer whales. *The Journal of the Acoustical Society of America*, 135(2), 975–993. <https://doi.org/10.1121/1.4861346>
- Miller, P. J. O., Johnson, M., & Tyack, P. L. (2004). Sperm whale behaviour indicates the use of echolocation click buzzes ‘Creaks’ in prey capture. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271(1554), 2239–2247. <https://doi.org/10.1098/rspb.2004.2863>
- Miller, P. J. O., Kvadsheim, P. H., Lam, F. P. A., Tyack, P. L., Curé, C., DeRuiter, S. L., Kleivane, L., Sivle, L. D., van IJsselmuide, S. P., Visser, F., Wensveen, P. J., von Benda-Beckmann, A. M., Martín López, L. M., Narazaki, T., & Hooker, S. K. (2015). First indications that northern bottlenose whales are sensitive to behavioural disturbance from anthropogenic noise. *Royal Society Open Science*, 2(6), 140484. <https://doi.org/10.1098/rsos.140484>
- Nagy, K. A. (2005). Field metabolic rate and body size. *Journal of Experimental Biology*, 208(9), 1621–1625. <https://doi.org/10.1242/jeb.01553>
- Nagy, K. A., Girard, I. A., & Brown, T. K. (1999). Energetics of free-ranging mammals, reptiles, and birds. *Annual Review of Nutrition*, 19(1), 247–277. <https://doi.org/10.1146/annurev.nutr.19.1.247>
- Nemoto, T. (1970). Feeding pattern of baleen whales in the ocean. In J. H. Steele (Ed.), *Marine Food Chains* (pp. 241–252). University of California Press.
- New, L. F., Moretti, D. J., Hooker, S. K., Costa, D. P., & Simmons, S. E. (2013). Using energetic models to investigate the survival and reproduction of beaked whales (Family Ziphiidae). *PLoS ONE*, 8(7), e68725. <https://doi.org/10.1371/journal.pone.0068725>
- Noren, D. P., Holt, M. M., Dunkin, R. C., Thometz, N. M., & Williams, T. M. (2016). Comparative and cumulative energetic costs of odontocete responses to anthropogenic disturbance. *Proceedings of Meetings on Acoustics*, 27(1), 040011. <https://doi.org/10.1121/2.0000357>
- Oli, M. K. (2004). The fast-slow continuum and mammalian life-history patterns: An empirical evaluation. *Basic and Applied Ecology*, 5(5), 449–463. <https://doi.org/10.1016/j.baae.2004.06.002>
- Pagel, M. D., Harvey, P. H., & Godfray, H. C. J. (1991). Species-abundance, biomass, and resource-use distributions. *The American Naturalist*, 138(4), 836–850. <https://doi.org/10.1086/285255>
- Parsons, E. C. M. (2017). Impacts of navy sonar on whales and dolphins: Now beyond a smoking gun? *Frontiers in Marine Science*, 4(295). <https://doi.org/10.3389/fmars.2017.00295>
- Pirotta, E., Booth, C. G., Cade, D. E., Calambokidis, J., Costa, D. P., Fahlbusch, J. A., Friedlaender, A. S., Goldbogen, J. A., Harwood, J., Hazen, E. L., New, L., & Southall, B. L. (2021). Context-dependent variability in the predicted daily energetic costs of disturbance for blue whales. *Conservation Physiology*, 9, coaa137. <https://doi.org/10.1093/conphys/coaa137>
- Pirotta, E., Booth, C. G., Costa, D. P., Fleishman, E., Kraus, S. D., Lusseau, D., Moretti, D., New, L. F., Schick, R. S., Schwarz, L. K., Simmons, S. E., Thomas, L., Tyack, P. L., Weise, M. J., Wells, R. S., & Harwood, J. (2018). Understanding the population consequences of disturbance. *Ecology and Evolution*, 8(19), 9934–9946. <https://doi.org/10.1002/ece3.4458>
- Popper, A. N., & Hawkins, A. D. (Eds.). (2012). *The effects of noise on aquatic life*. Springer.
- Potvin, J., Goldbogen, J. A., & Shadwick, R. E. (2012). Metabolic expenditures of lunge feeding rorquals across scale: Implications for the evolution of filter feeding and the limits to maximum body size. *PLoS ONE*, 7(9), e44854. <https://doi.org/10.1371/journal.pone.0044854>
- R Core Team. (2018). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.
- Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P.-O. (2010). Personality and the emergence of the pace-of-life syndrome concept at the population level. *Philosophical*

- Transactions of the Royal Society B: Biological Sciences*, 365(1560), 4051–4063. <https://doi.org/10.1098/rstb.2010.0208>
- Rohr, J. J., & Fish, F. E. (2004). Strouhal numbers and optimization of swimming by odontocete cetaceans. *Journal of Experimental Biology*, 207(10), 1633–1642. <https://doi.org/10.1242/jeb.00948>
- Rojano-Doñate, L., McDonald, B. I., Wisniewska, D. M., Johnson, M., Teilmann, J., Wahlberg, M., Højer-Kristensen, J., & Madsen, P. T. (2018). High field metabolic rates of wild harbour porpoises. *Journal of Experimental Biology*, 221(23), jeb185827. <https://doi.org/10.1242/jeb.185827>
- Sato, K., Takahashi, A., Miller, P. J. O., Tanaka, H., Kawabe, R., Ponganis, P. J., Handrich, Y., Akamatsu, T., Watanabe, Y., Mitani, Y., Costa, D. P., Bost, C.-A., Aoki, K., Amano, M., Trathan, P., Shapiro, A., & Naito, Y. (2007). Stroke frequency, but not swimming speed, is related to body size in free-ranging seabirds, pinnipeds and cetaceans. *Proceedings of the Royal Society B: Biological Sciences*, 274(1609), 471–477. <https://doi.org/10.1098/rspb.2006.0005>
- Schwacke, L. H., Smith, C. R., Townsend, F. I., Wells, R. S., Hart, L. B., Balmer, B. C., Collier, T. K., De Guise, S., Fry, M. M., Guillette, L. J., Lamb, S. V., Lane, S. M., McFee, W. E., Place, N. J., Tumlin, M. C., Ylitalo, G. M., Zolman, E. S., & Rowles, T. K. (2014). Health of common bottlenose dolphins (*Tursiops truncatus*) in Barataria Bay, Louisiana, Following the Deepwater Horizon Oil Spill. *Environmental Science & Technology*, 48(1), 93–103. <https://doi.org/10.1021/es403610f>
- Senigaglia, V., Christiansen, F., Bejder, L., Gendron, D., Lundquist, D., Noren, D. P., Schaffar, A., Smith, J. C., Williams, R., Martinez, E., Stockin, K., & Lusseau, D. (2016). Meta-analyses of whale-watching impact studies: Comparisons of cetacean responses to disturbance. *Marine Ecology Progress Series*, 542, 251–263. <https://doi.org/10.3354/meps11497>
- Shannon, G., McKenna, M. F., Angeloni, L. M., Crooks, K. R., Fristrup, K. M., Brown, E., Warner, K. A., Nelson, M. D., White, C., Briggs, J., McFarland, S., & Wittemyer, G. (2016). A synthesis of two decades of research documenting the effects of noise on wildlife. *Biological Reviews*, 91(4), 982–1005. <https://doi.org/10.1111/brv.12207>
- Shearer, J. M., Quick, N. J., Cioffi, W. R., Baird, R. W., Webster, D. L., Foley, H. J., Swaim, Z. T., Waples, D. M., Bell, J. T., & Read, A. J. (2019). Diving behaviour of Cuvier's beaked whales (*Ziphius cavirostris*) off Cape Hatteras, North Carolina. *Royal Society Open Science*, 6(2), 181728. <https://doi.org/10.1098/rsos.181728>
- Simpson, S. D., Radford, A. N., Nedelec, S. L., Ferrari, M. C. O., Chivers, D. P., McCormick, M. I., & Meekan, M. G. (2016). Anthropogenic noise increases fish mortality by predation. *Nature Communications*, 7, 10544. <https://doi.org/10.1038/ncomms10544>
- Sims, D. W., Southall, E. J., Humphries, N. E., Hays, G. C., Bradshaw, C. J. A., Pitchford, J. W., James, A., Ahmed, M. Z., Brierley, A. S., Hindell, M. A., Morritt, D., Musyl, M. K., Righton, D., Shepard, E. L. C., Wearmouth, V. J., Wilson, R. P., Witt, M. J., & Metcalfe, J. D. (2008). Scaling laws of marine predator search behaviour. *Nature*, 451(7182), 1098–1102. <https://doi.org/10.1038/nature06518>
- Sivle, L. D., Kvadsheim, P. H., Curé, C., Isojunno, S., Wensveen, P. J., Lam, F.-P., Visser, F., Kleivane, L., Tyack, P. L., Harris, C. M., & Miller, P. J. O. (2015). Severity of expert-identified behavioural responses of humpback whale, mink whale, and northern bottlenose whale to naval sonar. *Aquatic Mammals*, 41(4), 469–502. <https://doi.org/10.1578/AM.41.4.2015.469>
- Sivle, L. D., Wensveen, P. J., Kvadsheim, P. H., Lam, F.-P., Visser, F., Curé, C., Harris, C. M., Tyack, P. L., & Miller, P. J. O. (2016). Naval sonar disrupts foraging in humpback whales. *Marine Ecology Progress Series*, 562, 211–220. <https://doi.org/10.3354/meps11969>
- Sørensen, P. M., Wisniewska, D. M., Jensen, F. H., Johnson, M., Teilmann, J., & Madsen, P. T. (2018). Click communication in wild harbour porpoises (*Phocoena phocoena*). *Scientific Reports*, 8(1), 1–11. <https://doi.org/10.1038/s41598-018-28022-8>
- Southall, B. L., DeRuiter, S. L., Friedlaender, A., Stimpert, A. K., Goldbogen, J. A., Hazen, E., Casey, C., Fregosi, S., Cade, D. E., Allen, A. N., Harris, C. M., Schorr, G., Moretti, D., Guan, S., & Calambokidis, J. (2019). Behavioral responses of individual blue whales (*Balaenoptera musculus*) to mid-frequency military sonar. *Journal of Experimental Biology*, 222(5), jeb190637. <https://doi.org/10.1242/jeb.190637>
- Southall, B. L., Nowacek, D. P., Miller, P. J. O., & Tyack, P. L. (2016). Experimental field studies to measure behavioral responses of cetaceans to sonar. *Endangered Species Research*, 31, 293–315. <https://doi.org/10.3354/esr00764>
- Speakman, J. R. (2005). Body size, energy metabolism and lifespan. *Journal of Experimental Biology*, 208(9), 1717–1730. <https://doi.org/10.1242/jeb.01556>
- Stafford, K. M., Fox, C. G., & Clark, D. S. (1998). Long-range acoustic detection and localization of blue whale calls in the Northeast Pacific Ocean. *The Journal of the Acoustical Society of America*, 104(6), 3616–3625. <https://doi.org/10.1121/1.423944>
- Stahl, J. T., & Oli, M. K. (2006). Relative importance of avian life-history variables to population growth rate. *Ecological Modelling*, 198(1), 23–39. <https://doi.org/10.1016/j.ecolmodel.2006.04.001>
- Stimpert, A. K., DeRuiter, S. L., Southall, B. L., Moretti, D. J., Falcone, E. A., Goldbogen, J. A., Friedlaender, A., Schorr, G. S., & Calambokidis, J. (2014). Acoustic and foraging behavior of a Baird's beaked whale, *Berardius bairdii*, exposed to simulated sonar. *Scientific Reports*, 4, 7031. <https://doi.org/10.1038/srep07031>
- Sugihara, G. (1980). Minimal community structure: An explanation of species abundance patterns. *The American Naturalist*, 116(6), 770–787. <https://doi.org/10.1086/283669>
- Sumich, J. L. (1983). Swimming velocities, breathing patterns, and estimated costs of locomotion in migrating gray whales, *Eschrichtius robustus*. *Canadian Journal of Zoology*, 61(3), 647–652. <https://doi.org/10.1139/z83-086>
- Sutherland, W. J. (2007). Future directions in disturbance research. *Ibis*, 149(s1), 120–124. <https://doi.org/10.1111/j.1474-919X.2007.00673.x>
- Thompson, P. M., Lusseau, D., Barton, T., Simmons, D., Rusin, J., & Bailey, H. (2010). Assessing the responses of coastal cetaceans to the construction of offshore wind turbines. *Marine Pollution Bulletin*, 60(8), 1200–1208. <https://doi.org/10.1016/j.marpolbul.2010.03.030>
- Tollit, D., Harwood, J., Booth, C., Thomas, L., New, L., & Wood, J. (2016). *Cook Inlet Beluga Whale PCoD Expert Elicitation Workshop Report*. SMRUC-NA-NOAA0915. SMRU Consulting North America.
- Trites, A. W., & Pauly, D. (1998). Estimating mean body masses of marine mammals from maximum body lengths. *Canadian Journal of Zoology*, 76(5), 886–896. <https://doi.org/10.1139/z97-252>
- Tyack, P. L. (1986). Population biology, social behavior and communication in whales and dolphins. *Trends in Ecology & Evolution*, 1(6), 144–150. [https://doi.org/10.1016/0169-5347\(86\)90042-X](https://doi.org/10.1016/0169-5347(86)90042-X)
- Tyack, P. L., Zimmer, W. M. X., Moretti, D., Southall, B. L., Claridge, D. E., Durban, J. W., Clark, C. W., D'Amico, A., DiMarzio, N., Jarvis, S., McCarthy, E., Morrissey, R., Ward, J., & Boyd, I. L. (2011). Beaked whales respond to simulated and actual navy sonar. *PLoS ONE*, 6(3), e17009. <https://doi.org/10.1371/journal.pone.0017009>
- Villegas-Amtmann, S., Schwarz, L. K., Sumich, J. L., & Costa, D. P. (2015). A bioenergetics model to evaluate demographic consequences of disturbance in marine mammals applied to gray whales. *Ecosphere*, 6(10), art183. <https://doi.org/10.1890/ES15-00146.1>
- Watanabe, Y. Y., Goldman, K. J., Caselle, J. E., Chapman, D. D., & Papastamatiou, Y. P. (2015). Comparative analyses of animal-tracking data reveal ecological significance of endothermy in fishes. *Proceedings of the National Academy of Sciences of the United States of America*, 112(19), 6104–6109. <https://doi.org/10.1073/pnas.1500316112>
- Watanabe, Y. Y., Sato, K., Watanuki, Y., Takahashi, A., Mitani, Y., Amano, M., Aoki, K., Narazaki, T., Iwata, T., Minamikawa, S., & Miyazaki, N.

- (2011). Scaling of swim speed in breath-hold divers. *Journal of Animal Ecology*, 80(1), 57–68. <https://doi.org/10.1111/j.1365-2656.2010.01760.x>
- Watwood, S. L., Miller, P. J. O., Johnson, M., Madsen, P. T., & Tyack, P. L. (2006). Deep-diving foraging behaviour of sperm whales (*Physeter macrocephalus*). *Journal of Animal Ecology*, 75(3), 814–825. <https://doi.org/10.1111/j.1365-2656.2006.01101.x>
- Wensveen, P. J., Isojunno, S., Hansen, R. R., von Benda-Beckmann, A. M., Kleivane, L., van IJsselmuide, S., Lam, F.-P., Kvasdheim, P. H., DeRuiter, S. L., Curé, C., Narazaki, T., Tyack, P. L., & Miller, P. J. O. (2019). Northern bottlenose whales in a pristine environment respond strongly to close and distant navy sonar signals. *Proceedings of the Royal Society B: Biological Sciences*, 286(1899), 20182592. <https://doi.org/10.1098/rspb.2018.2592>
- White, C. R., & Seymour, R. S. (2003). Mammalian basal metabolic rate is proportional to body mass^{2/3}. *Proceedings of the National Academy of Sciences of the United States of America*, 100(7), 4046–4049. <https://doi.org/10.1073/pnas.0436428100>
- White, C. R., & Seymour, R. S. (2004). Does basal metabolic rate contain a useful signal? Mammalian BMR allometry and correlations with a selection of physiological, ecological, and life-history variables. *Physiological and Biochemical Zoology*, 77(6), 929–941. <https://doi.org/10.1086/425186>
- Williams, R., Lusseau, D., & Hammond, P. S. (2006). Estimating relative energetic costs of human disturbance to killer whales (*Orcinus orca*). *Biological Conservation*, 133(3), 301–311. <https://doi.org/10.1016/j.biocon.2006.06.010>
- Williams, T. M. (1999). The evolution of cost efficient swimming in marine mammals: Limits to energetic optimization. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 354(1380), 193–201. <https://doi.org/10.1098/rstb.1999.0371>
- Williams, T. M., Friedl, W. A., & Haun, J. E. (1993). The physiology of bottlenose dolphins (*Tursiops truncatus*): Heart rate, metabolic rate and plasma lactate concentration during exercise. *Journal of Experimental Biology*, 179(1), 31–46. <https://doi.org/10.1242/jeb.179.1.31>
- Williams, T. M., Kendall, T. L., Richter, B. P., Ribeiro-French, C. R., John, J. S., Odell, K. L., Losch, B. A., Feuerbach, D. A., & Andrew Stamper, M. (2017). Swimming and diving energetics in Dolphins: A stroke-by-stroke analysis for predicting the cost of flight responses in wild odontocetes. *Journal of Experimental Biology*, 220(6), 1135–1145. <https://doi.org/10.1242/jeb.154245>
- Wisniewska, D. M., Johnson, M., Teilmann, J., Rojano-Doñate, L., Shearer, J., Sveegaard, S., Miller, L. A., Siebert, U., & Madsen, P. T. (2016). Ultra-high foraging rates of harbor porpoises make them vulnerable to anthropogenic disturbance. *Current Biology*, 26(11), 1441–1446. <https://doi.org/10.1016/j.cub.2016.03.069>
- Wisniewska, D. M., Johnson, M., Teilmann, J., Siebert, U., Galatius, A., Dietz, R., & Madsen, P. T. (Eds.). (2018). High rates of vessel noise disrupt foraging in wild harbour porpoises (*Phocoena phocoena*). *Proceedings of the Royal Society B: Biological Sciences*. 285(1872), 20172314. <https://doi.org/10.1098/rspb.2017.2314>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Czapaniskiy MF, Savoca MS, Gough WT, et al. Modelling short-term energetic costs of sonar disturbance to cetaceans using high-resolution foraging data. *J Appl Ecol*. 2021;00:1–15. <https://doi.org/10.1111/1365-2664.13903>